

Situational awareness and forecasting for Norway

FHI COVID-19 modelling team

Week 35, 14 September 2022

Main results:

We report the estimated national reproduction number from our changepoint model this week. The model estimates an average effective reproduction number of around 0.78 since 11 July. Due to a lack of reliable test data, we only base our estimates on hospitalisation data, which has a longer delay. We also refer to trend analysis results Figure 13 based on various data sources.

- **National epidemiological situation:**

The most recent reproduction numbers:

Model	Median	2.5%	97.5%	Prob>1	period/day	More info
Changepoint	0.78	0.71	0.85	$\leq 1\%$	from Jul 11	Section 1

- **National forecasting:**

One-week-ahead national forecasts from changepoint model:

Indicator	Median/Mean	95% PI	day	Info 2-3 weeks forecasts
Hospital beds	53/52	(35-75)	Sep 18	Table 2
Ventilator beds	2/1	(0-4)	Sep 18	Table 2

Three-weeks-ahead national forecasts can be found in Tables 2. Age-specific hospital prevalence predictions are provided in Figures 4 and 5.

- **Regional epidemiological situation:**

The newest regional effective reproduction number for Oslo:

Model	Median	2.5%	97.5%	Prob>1	day	Info other counties
SMC	0.34	0.15	0.63	≤ 0.01	24 Aug - 27 Aug	Table 3

- **Telenor mobility data and the number of foreign visitors:**

Since 1 July 2022 we no longer receive real-time mobility data from Telenor.

1 Estimated national reproduction numbers

Table 1 shows the estimated reproductive number of our national changepoint model. Figure 1 shows the estimated daily number of COVID-19 patients admitted to hospital, with blue medians and interquartile bands, which are compared to the actual true data, provided in red. The uncertainty captures the uncertainty in the calibrated parameters in addition to the stochastic elements of our model and the variability of other model parameters.

Table 1: Calibration results

Reff	Period
2.7/2.72(2.3-3.17)	From Feb 17 to Mar 14
0.69/0.69(0.57-0.8)	From Mar 15 to Apr 19
0.41/0.43(0.06-0.91)	From Apr 20 to May 10
0.85/0.79(0.17-1.25)	From May 11 to Jun 30
0.53/0.58(0.07-1.3)	From Jul 01 to Jul 31
1.15/1.13(0.59-1.53)	From Aug 01 to Aug 31
0.98/0.99(0.81-1.18)	From Sep 01 to Oct 25
1.24/1.25(0.77-1.74)	From Oct 26 to Nov 04
0.99/0.99(0.85-1.14)	From Nov 05 to Nov 30
0.81/0.81(0.7-0.94)	From Dec 01 to Jan 03
0.8/0.8(0.52-1.04)	From Jan 04 to Jan 21
0.87/0.88(0.6-1.22)	From Jan 22 to Feb 07
1.29/1.3(1.09-1.53)	From Feb 08 to Mar 01
1/1(0.89-1.11)	From Mar 02 to Mar 24
0.85/0.84(0.77-0.91)	From Mar 25 to May 05
0.85/0.85(0.62-1.11)	From May 06 to May 26
0.69/0.69(0.42-0.95)	From May 27 to Jun 20
0.9/0.87(0.5-1.12)	From Jun 21 to Aug 04
1.05/1.05(0.87-1.28)	From Aug 05 to Aug 31
0.77/0.77(0.65-0.88)	From Sep 01 to Sep 24
1.03/1.03(0.99-1.08)	From Sep 25 to Dec 14
0.97/0.97(0.89-1.05)	From Dec 15 to Jan 13
1.21/1.2(1.14-1.26)	From Jan 14 to Feb 25
0.7/0.7(0.66-0.74)	From Feb 26 to May 14
1.06/1.06(0.98-1.13)	From May 15 to Jul 10
0.78/0.78(0.71-0.85)	From Jul 11
Median/Mean (95% credible intervals)	




Figure 1: A comparison of true data (red) and predicted values (blue) for hospital admissions . The last four data points (black) are assumed to be affected by reporting delay. The uncertainty captures the uncertainty in the calibrated parameters, in addition to the stochastic elements of our model and the variability of other model parameters.

In figure 2, we show how our national model fits the national hospital prevalence data (2a) and the daily number of patients receiving ventilator treatment (2b). Those data sources are not used to estimate the parameters, and can therefore be seen as a validation of the model assumptions.




Figure 2: A comparison of true data (red) and predicted values (blue) for hospital and respirator prevalence. Prevalence data is based on NIPaR and may be different to the data from Helsedirektoratet.

2 National 3-week predictions: Hospital beds and Ventilator bed

In this section we show the projected national prevalence of hospitalised patients (hospital beds) and prevalence of patients receiving ventilator treatment (ventilator beds). The projected epidemic and healthcare burden are illustrated in table 2.

Age-specific hospital prevalence predictions are provided in Figures 4 and 5.

Table 2: Estimated national hospital beds and ventilator beds. Median/Mean (CI)

	1 week prediction (Sep 18)	2 week prediction (Sep 25)	3 week prediction (Oct 02)
Hospital beds	53/52 (35-75)	50/50 (33-71)	48/47 (31-67)
Ventilator beds	2/1 (0-4)	2/1 (0-4)	1/1 (0-4)




Figure 3: National 3 week predictions for hospital beds (left) and ventilator beds (right)




Figure 4: Simulated hospital prevalence by age group taking into account the omicron takeover. Real data is shown as black dots.




Figure 5: Simulated respirator prevalence by age group taking into account the omicron takeover. Real data is shown as black dots.

3 Estimated regional reproduction numbers

Calibration of our regional SMC model to hospitalisation incidence data leads to the following estimates for current regional effective reproduction numbers by county (Table 3).

Below we show the estimated daily number of COVID-19 patients admitted to hospital in each county. Model estimates are shown with blue medians and interquartile bands, which are compared to the actual true data, provided in red. The blue bands describe the uncertainty in the calibrated parameters, in addition to the stochastic elements of our model. Last four data points are shown in black as they may be affected by reporting delay.

Table 3: Regional estimates, 24 Aug-27 Aug

County	Median	95CI	Prob>1
Oslo	0.34	0.15-0.63	0
Rogaland	0.44	0.18-0.87	0.01
Møre og Romsdal	0.54	0.28-0.92	0.01
Nordland	0.39	0.16-0.77	0
Viken	0.63	0.4-1.04	0.03
Innlandet	0.41	0.18-0.78	0
Vestfold og Telemark	0.70	0.41-1.24	0.1
Agder	0.59	0.28-1.15	0.06
Vestland	0.33	0.15-0.64	0
Trøndelag	0.50	0.26-0.84	0.01
Troms og Finnmark	0.54	0.25-1.08	0.04










Figure 6: The map shows the direction of the trend in incidence in the counties based on the latest effective reproduction numbers shown in the other chart. The trend is increasing if the probability that the latest reproduction number is above one is above 95%, the trend is likely increasing if this probability is between 80% and 95%, the trend is uncertain if the probability is between 20% and 80%, the trend is likely decreasing if the probability is between 5% and 20% and is decreasing if the probability that the latest R is above one is less than 5%.

Estimated vs observed hospital incidence and 3 weeks forecast by county:

Forecasts are now based on the estimated reproduction numbers obtained by our regional SMC model, for each county. In the forecasted period of three weeks, we use the reproductive numbers showed on Table 3.





4 Regional 3-week predictions: Cumulative (total) incidence and Prevalence

Below is shown the daily incidence (table 4) and prevalence (table 5) for each county.

Table 4: Predicted incidence per day: Median/Mean (CI)

Region	1 week prediction (19 Sep)	2 weeks prediction (26 Sep)	3 weeks prediction (03 Oct)
Agder	15/34 (2-147)	13/38 (2-160)	13/41 (2-168)
Innlandet	17/21 (5-62)	15/19 (4-59)	14/18 (4-56)
Møre og Romsdal	13/20 (2-77)	10/17 (1-70)	9/15 (1-65)
Nordland	4/6 (0-22)	4/6 (0-20)	4/5 (0-19)
Oslo	68/76 (27-173)	66/75 (26-175)	64/75 (26-181)
Rogaland	11/18 (2-68)	11/18 (2-70)	11/19 (2-76)
Troms og Finnmark	6/11 (0-46)	5/11 (0-48)	5/11 (0-47)
Trøndelag	25/36 (7-122)	22/32 (6-105)	20/30 (6-99)
Vestfold og Telemark	49/81 (13-366)	45/85 (11-429)	42/89 (11-497)
Vestland	17/24 (4-80)	16/22 (4-72)	15/22 (4-70)
Viken	172/243 (61-875)	159/242 (57-964)	151/242 (56-1008)

Table 5: Predicted prevalence. Number of infectious individuals (asymptomatic plus pre-symptomatic plus symptomatic) per day. Median/Mean and 95 perc. CI for three weeks prediction.

Region	19 Sep	26 Sep	03 Oct	low CI, 03 Oct	high CI, 03 Oct
Agder	95/172	85/186	79/202	35	742
Innlandet	99/116	88/105	82/99	42	262
Møre og Romsdal	86/117	72/101	65/93	32	305
Nordland	45/52	43/49	42/48	25	104
Oslo	756/790	745/783	741/782	564	1240
Rogaland	90/116	87/117	85/121	47	358
Troms og Finnmark	43/63	40/62	38/63	18	215
Trøndelag	173/217	154/196	144/185	82	498
Vestfold og Telemark	239/377	219/389	208/407	73	2105
Vestland	156/187	147/176	143/174	95	388
Viken	1061/1361	995/1346	955/1348	539	4713

5 Regional 3-week predictions: Hospital beds and ventilator beds

Below is shown the estimated short-term forecasting of expected hospital prevalence (table 6) and patients on ventilator treatment for each county (table 7). These estimates do not take into account imported cases from abroad that may be admitted to hospital.

Table 6: Number of hospitalisation beds occupied by Covid-19 patients: Median/Mean (CI)

Region	1 week prediction (19 Sep)	2 weeks prediction (26 Sep)	3 weeks prediction (03 Oct)
Agder	1/2 (0-10)	1/2 (0-12)	1/2 (0-12)
Innlandet	1/2 (0-7)	1/2 (0-7)	1/1 (0-7)
Møre og Romsdal	1/2 (0-8)	1/1 (0-7)	1/1 (0-7)
Nordland	0/1 (0-3)	0/0 (0-3)	0/0 (0-2)
Oslo	3/4 (0-12)	3/4 (0-12)	3/4 (0-11)
Rogaland	1/1 (0-6)	0/1 (0-6)	0/1 (0-6)
Troms og Finnmark	0/1 (0-5)	0/1 (0-4)	0/1 (0-3)
Trøndelag	2/3 (0-11)	2/2 (0-9)	1/2 (0-9)
Vestfold og Telemark	4/6 (0-24)	3/6 (0-28)	3/6 (0-33)
Vestland	1/2 (0-8)	1/2 (0-7)	1/1 (0-6)
Viken	12/16 (3-56)	10/16 (2-59)	10/15 (2-63)

Table 7: Number of ICU beds occupied by Covid-19 patients: Median/Mean (CI)

Region	1 week prediction (19 Sep)	2 weeks prediction (26 Sep)	3 weeks prediction (03 Oct)
Agder	0/0 (0-1)	0/0 (0-1)	0/0 (0-1)
Innlandet	0/0 (0-1)	0/0 (0-1)	0/0 (0-1)
Møre og Romsdal	0/0 (0-1)	0/0 (0-1)	0/0 (0-1)
Nordland	0/0 (0-0)	0/0 (0-0)	0/0 (0-0)
Oslo	0/0 (0-1)	0/0 (0-1)	0/0 (0-1)
Rogaland	0/0 (0-1)	0/0 (0-1)	0/0 (0-1)
Troms og Finnmark	0/0 (0-1)	0/0 (0-0)	0/0 (0-0)
Trøndelag	0/0 (0-1)	0/0 (0-1)	0/0 (0-1)
Vestfold og Telemark	0/0 (0-1)	0/0 (0-1)	0/0 (0-2)
Vestland	0/0 (0-1)	0/0 (0-1)	0/0 (0-1)
Viken	0/1 (0-3)	0/1 (0-3)	0/1 (0-3)

6 14-day trend analysis of confirmed cases and hospitalisations by county


To estimate recent trends in hospitalisation and number of positive tests, we present results in table 8 based on a negative binomial regression where we account for weekend effects. We exclude the last three days to avoid problems of reporting delay and fit the model using data from 17 days to 3 days before the current date. We fit a separate trend model for confirmed cases and for hospital incidence. We only fit a trend model if there has been more than 5 cases or hospitalisations in the 14-day period.

Table 8: Trend analysis for the last 14 days


County	Average daily increase last 14 days		Doubling Time (days)	
	Hospitalisations	Cases	Hospitalisations	Cases
Agder	-13.7 (-31.9, 5.1) %	-6.2 (-12.7, 0.6) %	-4.7 (-1.8, 14)	-10.8 (-5.1, 125.4)
Innlandet	Not enough data	-0.6 (-6.2, 5.3) %	Not enough data	-115.5 (-10.9, 13.5)
Møre og Romsdal	-10.7 (-27.5, 6.9) %	-12.9 (-18, -7.7) %	-6.1 (-2.2, 10.4)	-5 (-3.5, -8.7)
Nordland	Not enough data	-1.2 (-8.2, 6.2) %	Not enough data	-56.7 (-8.1, 11.5)
Norge	-5.4 (-9.3, -1.4) %	-3.9 (-5.7, -2) %	-12.5 (-7.1, -49.1)	-17.7 (-11.8, -34.9)
Oslo	-18.6 (-38.7, 2.2) %	-4.7 (-10.4, 1.3) %	-3.4 (-1.4, 31.8)	-14.4 (-6.3, 54.8)
Rogaland	Not enough data	-3.8 (-9.9, 2.7) %	Not enough data	-18.1 (-6.6, 26.1)
Troms og Finnmark	Not enough data	-0.5 (-7.3, 6.8) %	Not enough data	-134.6 (-9.1, 10.6)
Trøndelag	-0.5 (-14.2, 15.2) %	0.2 (-5, 5.6) %	-127.1 (-4.5, 4.9)	373.6 (-13.6, 12.7)
Vestfold og Telemark	-8.8 (-19.2, 2) %	-8.1 (-14.2, -1.9) %	-7.5 (-3.2, 35.3)	-8.2 (-4.5, -35.8)
Vestland	4.9 (-5.2, 16.6) %	-2.7 (-6.3, 1.1) %	14.4 (-13, 4.5)	-25.8 (-10.7, 64.6)
Viken	-7.2 (-14.3, 0.2) %	-3.1 (-8.1, 2.2) %	-9.3 (-4.5, 459.5)	-22.2 (-8.2, 31.3)

7 Regional SMC-model: Estimated daily reproduction numbers


In the figures below we plot the 95% credibility interval and quantiles of the estimated posterior distribution of the regional, daily reproduction numbers. For some counties, uncertainty is large towards the most recent time, because there are very few data and possibly reporting delays which are different in each county.




Vestfold og Telemark




Agder




Rogaland




Vestland






Innlandet



Møre og Romsdal





8 Short-term predictions per health trust ("helseforetak"): number of occupied covid-19 beds and number of patients receiving ventilator treatment

In the table below we distribute our short-term predictions to the Norwegian health trusts ("helseforetak"), on the basis of their areas of responsibility.

Table 9: Number of hospitalisation beds occupied by Covid-19 patients: Median/Mean (CI)


Helseforetak	1 week prediction (19 Sep)	2 weeks prediction (26 Sep)	3 weeks prediction (03 Oct)
Akershus universitetssykehus	5/6 (2-20)	4/6 (1-21)	4/6 (1-23)
Finnmarksykehuset	0/0 (0-1)	0/0 (0-1)	0/0 (0-1)
Helse Bergen	1/1 (0-6)	1/1 (0-4)	1/1 (0-4)
Helse Fonna	0/0 (0-2)	0/0 (0-1)	0/0 (0-2)
Helse Førde	0/0 (0-1)	0/0 (0-1)	0/0 (0-1)
Helse Møre og Romsdal	1/2 (0-8)	1/1 (0-7)	1/1 (0-6)
Helse Nord Trøndelag	1/1 (0-3)	1/1 (0-3)	0/1 (0-3)
Helse Stavanger	1/1 (0-4)	1/1 (0-4)	0/1 (0-4)
Nordlandssykehuset	0/0 (0-2)	0/0 (0-2)	0/0 (0-1)
Oslo universitetssykehus	3/3 (0-9)	3/3 (0-9)	3/3 (0-8)
St. Olavs hospital	1/2 (0-7)	1/2 (0-6)	1/1 (0-5)
Sykehuset Innlandet	1/2 (0-6)	1/2 (0-5)	1/1 (0-5)
Sykehuset Telemark	2/2 (0-10)	1/2 (0-11)	1/3 (0-13)
Sykehuset i Vestfold	2/3 (0-14)	2/3 (0-16)	2/4 (0-19)
Sykehuset Østfold	3/4 (1-14)	3/4 (1-15)	3/4 (1-16)
Sørlandet sykehus	1/2 (0-10)	1/2 (0-11)	1/2 (0-11)
Universitetssykehuset i Nord-Norge	0/1 (0-3)	0/1 (0-3)	0/1 (0-3)
Vestre Viken	5/6 (1-21)	4/6 (1-23)	4/6 (1-25)

Table 10: Number of respirator beds occupied by Covid-19 patients: Median/Mean (CI)


Helseforetak	1 week prediction (19 Sep)	2 weeks prediction (26 Sep)	3 weeks prediction (03 Oct)
Akershus universitetssykehus	0/0 (0-1)	0/0 (0-1)	0/0 (0-1)
Finnmarksykehuset	0/0 (0-0)	0/0 (0-0)	0/0 (0-0)
Helgelandssykehuset	0/0 (0-0)	0/0 (0-0)	0/0 (0-0)
Helse Bergen	0/0 (0-1)	0/0 (0-1)	0/0 (0-1)
Helse Fonna	0/0 (0-0)	0/0 (0-0)	0/0 (0-0)
Helse Førde	0/0 (0-0)	0/0 (0-0)	0/0 (0-0)
Helse Møre og Romsdal	0/0 (0-1)	0/0 (0-1)	0/0 (0-1)
Helse Nord Trøndelag	0/0 (0-0)	0/0 (0-0)	0/0 (0-0)
Helse Stavanger	0/0 (0-1)	0/0 (0-1)	0/0 (0-1)
Nordlandssykehuset	0/0 (0-0)	0/0 (0-0)	0/0 (0-0)
Oslo universitetssykehus	0/0 (0-1)	0/0 (0-1)	0/0 (0-1)
St. Olavs hospital	0/0 (0-1)	0/0 (0-1)	0/0 (0-1)
Sykehuset Innlandet	0/0 (0-1)	0/0 (0-1)	0/0 (0-1)
Sykehuset Telemark	0/0 (0-0)	0/0 (0-0)	0/0 (0-1)
Sykehuset i Vestfold	0/0 (0-1)	0/0 (0-1)	0/0 (0-1)
Sykehuset Østfold	0/0 (0-1)	0/0 (0-1)	0/0 (0-1)
Sørlandet sykehus	0/0 (0-1)	0/0 (0-1)	0/0 (0-1)
Universitetssykehuset i Nord-Norge	0/0 (0-1)	0/0 (0-0)	0/0 (0-0)
Vestre Viken	0/0 (0-1)	0/0 (0-1)	0/0 (0-1)

In the figures below, we plot the 95% credibility interval and quantiles of the estimated posterior distribution of the daily number of occupied covid-19 beds and the number of patients receiving ventilator treatment for a few health trusts. The others are available on request. In blue are actual data, in green our median prediction and in gray the 95% credibility interval. For some counties, uncertainty is considerable towards the most recent time because there is little data and possibly reporting delays that are different in each county.


Akershus universitetssykehus




Akershus universitetssykehus




Oslo universitetssykehus




Oslo universitetssykehus



St. Olavs hospital



St. Olavs hospital



9 Estimated parameters

Table 11: Estimated parameters

	Min.	1st Qu.	Median	Mean	3rd Qu.	Max.	Period
R0s	2.083	2.573	2.705	2.72	2.865	3.316	Until March 14
R1s	0.45	0.65	0.687	0.687	0.728	0.873	From 15 March to 19 April
R2s	0.006	0.288	0.406	0.433	0.572	1.155	From 20 April to 10 May
R3s	0.056	0.602	0.845	0.788	1.006	1.378	From 11 May to 30 June
R4s	0	0.33	0.528	0.582	0.813	1.712	From 01 July to 31 July
R5s	0.386	0.974	1.149	1.131	1.311	1.67	From 01 August to 31 August
R6s	0.678	0.926	0.985	0.988	1.043	1.271	From 01 September to 25 October
R7s	0.521	1.095	1.238	1.249	1.404	2.003	From 26 October to 04 November
R8s	0.752	0.933	0.992	0.989	1.043	1.207	From 05 November to 30 November
R9s	0.633	0.766	0.805	0.809	0.847	1.028	From 01 December to 03 January
R10s	0.416	0.706	0.797	0.799	0.9	1.192	From 04 January to 21 January
R11s	0.468	0.771	0.872	0.882	0.994	1.377	From 22 January to 07 February
R12s	1.038	1.212	1.292	1.298	1.384	1.648	From 08 February to 01 March
R13s	0.834	0.956	0.997	0.997	1.035	1.178	From 02 March to 24 March
R14s	0.727	0.816	0.846	0.841	0.864	0.939	From 25 March to 05 May
R15s	0.517	0.778	0.849	0.855	0.932	1.258	From 06 May to 26 May
R16s	0.147	0.605	0.693	0.689	0.78	1.09	From 27 May to 20 June
R17s	0.345	0.788	0.896	0.874	0.976	1.212	From 21 June to 04 August
R18s	0.794	0.984	1.046	1.053	1.11	1.422	From 05 August to 31 August
R19s	0.59	0.729	0.771	0.768	0.806	0.936	From 01 September to 24 September
R20s	0.946	1.016	1.034	1.034	1.053	1.102	From 25 September to 14 December
R21s	0.855	0.943	0.97	0.971	0.997	1.104	From 15 December to 13 January
R22s	1.101	1.184	1.208	1.204	1.226	1.286	From 14 January to 25 February
R23s	0.63	0.683	0.697	0.698	0.711	0.774	From 26 February to 14 May
R24s	0.942	1.039	1.062	1.06	1.085	1.154	From 15 May to 10 July
R25s	0.652	0.759	0.782	0.781	0.806	0.89	From 11 July

Table 12: Assumptions

Assumptions	Mean	Distribution	Reference
Mobile Mobility Data			
Telenor coverage	48%		https://ekomstatistikken.nkom.no/
Data updated	Jun 5		
Data used in the predictions	Jun 3 rd	Fixed	Corrected to preserve population
Model parameters			
Exposed period ($1/\lambda_1$)	2 days	Exponential	changed from Feretti et al 2020
Pre-symptomatic period ($1/\lambda_2$)	2 days	Exponential	Feretti et al 2020
Symptomatic infectious period ($1/\gamma$)	3 days	Exponential	changed from Feretti et al 2020
Asymptomatic, infectious period ($1/\gamma$)	3 days	Exponential	changed from Feretti et al 2020
Infectiousness asympt. (r_{I_a})	0.1	Fixed	Feretti et al 2020
Infectiousness presymp (r_{E_2})	1.3	Fixed	guided by Feretti et al 2020
Prob. asymptomatic infection (p_a)	0.4		Feretti et al 2020
Healthcare			
Fraction asymptomatic infections	40%	Fixed	Mizumoto et al 2020 20% for the old population, Diamond Princess
% symptomatic and asymptomatic infections requiring hospitalization:			Salje et al 2020 corrected for: % of elderly living in elderly homes in Norway (last two age groups) and corrected for presence among positive tested since May 1.
0-9 years	0.1%	Fixed	
10 - 19 years	0.1%		
20 - 29 years	0.5%		
30 - 39 years	1.1%		
40 - 49 years	1.4%		
50 - 59 years	2.9%		
60 - 69 years	5.8%		
70 - 79 years	9.3%		
80+ years	22.3%		
Probability that an admission has been reported on Monday			
From Sunday	32%	Fixed	Estimated from "Beredskapsregistret BeredtC19"
From Saturday	49%		
From Friday	68%		
From Thursday	86%		
Probability that an admission has been reported			
From one day before	53%	Fixed	Estimated from "Beredskapsregistret BeredtC19"
From two days before	77%		
From three days before	82%		
From four days before	91%		
Probability that a positive laboratory test has been reported			
From one day before	6.7%	Fixed	Estimated from MSIS
From two days before	59%		
From three days before	90%		
From four days before	97%		
Probability that a negative laboratory test has been reported			
From one day before	16%	Fixed	Estimated from MSIS
From two days before	74%		
From three days before	92%		
From four days before	98%		

Supplementary analysis: Trend analysis using various data sources

Due to significant changes in test recommendations, we present trend calculations from several monitoring data sources. Seen together, they can give insights on the trend of new infections. The data sources include:

- MSIS - A trend calculated from the number of confirmed positive cases. Changes in test criteria in recent weeks will likely lead to underestimating the trend of new infections.
- NOPaR: Admissions with covid-19 as the main reason, the trend is calculated from daily numbers admissions with covid-19 as the main cause. It usually takes longer from infection to hospitalization than for testing; therefore, this indicator lags behind the trend of transmission at around 1-2 weeks. The major difference in severity between Delta and Omicron also makes this indicator more challenging to interpret in the transition phase between the two variants.
- Symptometer - From the Symptometer survey, we calculate the proportion of respondents who self-report symptoms and a positive test for covid-19. This indicator provides a measure of the prevalence of infection in society, but after conversion, it can also provide an estimate of a trend for new infections.
- sKUHR - We use the development of the number of consultations with confirmed covid-19 (R992) at GP offices and emergency rooms, adjusting for delayed reporting the last 14 days. After correction for the delay between onset and seeking a doctor, this indicator measures the incidence of infection.
- MoBa - The participants in MoBa are sent a mobile questionnaire form every other week, including questions about who has been "sick with respiratory symptoms/fever the last 14 days" and how many days ago the symptoms started. We calculate an approximate 14-day incidence of symptom onset from this indicator, which is used to estimate a trend in infection. The number of respondents is about 60-75,000 in each round.
- NPR and NoPaR: Proportion of all acute admissions with covid-19, including entries with other main causes than covid-19. This indicator measures the prevalence of infection in society and is then converted into incidence of infection.




Figure 13: Trend analysis using various data sources




Figure 14: Regional hospitalisation probabilities per infection. The estimates are based on Salje et al., and regional data on the age distribution in the test data and the empirical case-hospitalisation rates.

Models and materials:

This report presents results based on a mathematical infectious disease model describing the geographical spread of COVID-19 in Norway. We use a metapopulation model for situational awareness and short-term forecasting and an individual-based model for long-term predictions. This report does not contain the long-term prediction results. Reproduction numbers of the metapopulation model are estimated in two ways: SMC-ABC is used to estimate a step-function in the transmissibility through prespecified changepoints, and SMC is used to estimate a daily varying reproduction number. We also provide estimates based on EpiEstim and a simple trend analysis. The models are described in previous reports and will not be explained here.

The metapopulation model takes daily varying Telenor mobility data as input. We also provide plots of the recent mobility for situational awareness.

How you should interpret the results: 3-week-ahead predictions and long-term scenarios

We provide both 3-week-ahead predictions and long-term scenarios. These are simulations of the disease spread into the future, under specific assumptions.

In the 3-week-ahead predictions, we assume that all parameters are as today, and simulate disease spread 3-weeks-ahead in time. Hence, these predictions are conditional on the current situation, and specifically on the most recently estimated reproduction number. The 3-week-ahead predictions thus do not take into account changes in transmissibility that are not yet captured by the available data, for example due to the delay between transmission and hospitalisation. Hence one of the conditions for the predictions to be valid is that the intervention policies do not change significantly in the next weeks. Hence, it does not make sense to evaluate or use the predictions if there are big changes in factors like

- new interventions
- relaxation of interventions
- a combination of new interventions and relaxations
- a significant change in vaccination coverage
- new variants with new properties
- a significant change in the contact behaviour of individuals.

As these factors are not likely to stay constant in the long-term future, we do not produce predictions for longer than three weeks ahead in time. Hence, our 3-week-ahead predictions are predictions of what may happen in the future, if there were no significant changes in the assumptions.

In this report, the term patient in ventilator treatment includes only those patients that require either invasive mechanical ventilation or ECMO (Extracorporeal membrane oxygenation).

FHI COVID-19 modelling team:

- **Birgitte Freiesleben de Blasio** - Department of Method Development and Analytics. Norwegian Institute of Public Health and Oslo Centre for Biostatistics and Epidemiology, University of Oslo.
- **Francesco Di Ruscio** - Department of Method Development and Analytics. Norwegian Institute of Public Health.
- **Gunnar Øyvind Isaksson Rø** - Department of Method Development and Analytics. Norwegian Institute of Public Health.
- **Solveig Engebretsen** - Norsk Regnesentral.
- **Arnoldo Frigessi** - Oslo Centre for Biostatistics and Epidemiology, University of Oslo and Oslo University Hospital.
- **Alfonso Diz-Lois Palomares** - Department of Method Development and Analytics. Norwegian Institute of Public Health.
- **Magnus Nygård Osnes** - Department of Method Development and Analytics. Norwegian Institute of Public Health.
- **Anja Bråthen Kristoffersen** - Department of Method Development and Analytics. Norwegian Institute of Public Health.
- **Kenth Engø-Monsen** - Telenor Research.
- **Louis Yat Hin Chan** - Department of Method Development and Analytics. Norwegian Institute of Public Health.
- **Jonas Christoffer Lindstrøm** - Department of Method Development and Analytics. Norwegian Institute of Public Health.
- **Richard White** - Department of Method Development and Analytics. Norwegian Institute of Public Health.
- **Gry Marysol Grøneng** - Department of Method Development and Analytics. Norwegian Institute of Public Health.
- **Chi Zhang** - Department of Method Development and Analytics. Norwegian Institute of Public Health.
- **Jørgen Eriksson Midtbø** - Department of Method Development and Analytics. Norwegian Institute of Public Health.
- **Geir Storvik** - Department of Mathematics. University of Oslo.