

Situational awareness and forecasting for Norway

FHI COVID-19 modelling team

Week 25, 22 June 2022

Main results:

We report the estimated national reproduction number from our changepoint model this week. The model estimates an average effective reproduction number of around 1.2 since 16 May. Due to a lack of reliable test data, we only base our estimates on hospitalisation data, which has a longer delay. We also refer to trend analysis results Figure 21 based on various data sources.

- **National epidemiological situation:**

The most recent reproduction numbers:

Model	Median	2.5%	97.5%	Prob>1	period/day	More info
Changepoint	1.23	0.93	1.46	35%	from May 16	Section 1

- **National forecasting:**

One-week-ahead national forecasts from changepoint model:

Indicator	Median/Mean	95% PI	day	Info 2-3 weeks forecasts
Hospital beds	101/96	(47-187)	Jun 26	Table 2
Ventilator beds	5/4	(1-10)	Jun 26	Table 2

Three-weeks-ahead national forecasts can be found in Tables 2. Age-specific hospital prevalence predictions are provided in Figures 4 and 5.

- **Regional epidemiological situation:**

The newest regional effective reproduction number for Oslo:

Model	Median	2.5%	97.5%	Prob>1	day	Info other counties
SMC	0.52	0.29	0.88	0.01	01 Jun - 04 Jun	Table 3

- **Telenor mobility data and the number of foreign visitors:** The most recent mobility development is found in Figures 15 and 16 for the largest municipality in each county and for each county, respectively. Figure 17 shows the total number of roamers per day per county. Overall, there has been an increasing volatility in the inter-county mobility since Easter.

1 Estimated national reproduction numbers

Table 1 shows the estimated reproductive number of our national changepoint model. Figure 1 shows the estimated daily number of COVID-19 patients admitted to hospital, with blue medians and interquartile bands, which are compared to the actual true data, provided in red. The uncertainty captures the uncertainty in the calibrated parameters in addition to the stochastic elements of our model and the variability of other model parameters.

Table 1: Calibration results

Reff	Period
2.69/2.7(2.19-3.25)	From Feb 17 to Mar 14
0.68/0.68(0.55-0.81)	From Mar 15 to Apr 19
0.49/0.49(0.03-0.99)	From Apr 20 to May 10
0.53/0.54(0.04-1.25)	From May 11 to Jun 30
0.56/0.58(0.05-1.3)	From Jul 01 to Jul 31
1.21/1.2(0.74-1.56)	From Aug 01 to Aug 31
1.01/1(0.83-1.17)	From Sep 01 to Oct 25
1.11/1.13(0.67-1.74)	From Oct 26 to Nov 04
0.98/0.98(0.83-1.12)	From Nov 05 to Nov 30
0.86/0.85(0.72-0.97)	From Dec 01 to Jan 03
0.78/0.78(0.46-1.08)	From Jan 04 to Jan 21
0.86/0.88(0.57-1.26)	From Jan 22 to Feb 07
1.29/1.29(1.04-1.53)	From Feb 08 to Mar 01
1.01/1.01(0.88-1.14)	From Mar 02 to Mar 24
0.84/0.84(0.76-0.91)	From Mar 25 to May 05
0.89/0.88(0.65-1.09)	From May 06 to May 26
0.73/0.73(0.45-1.03)	From May 27 to Jun 20
0.8/0.78(0.41-1.1)	From Jun 21 to Aug 04
1.09/1.09(0.9-1.27)	From Aug 05 to Aug 31
0.75/0.75(0.6-0.9)	From Sep 01 to Sep 24
1.02/1.02(0.95-1.08)	From Sep 25 to Dec 14
1.07/1.07(0.97-1.16)	From Dec 15 to Jan 13
1.34/1.34(1.28-1.41)	From Jan 14 to Feb 25
0.93/0.93(0.87-0.98)	From Feb 26 to May 15
1.23/1.22(0.93-1.46)	From May 16
Median/Mean (95% credible intervals)	




Figure 1: A comparison of true data (red) and predicted values (blue) for hospital admissions . The last four data points (black) are assumed to be affected by reporting delay. The uncertainty captures the uncertainty in the calibrated parameters, in addition to the stochastic elements of our model and the variability of other model parameters.

In figure 2, we show how our national model fits the national hospital prevalence data (2a) and the daily number of patients receiving ventilator treatment (2b). Those data sources are not used to estimate the parameters, and can therefore be seen as a validation of the model assumptions.




Figure 2: A comparison of true data (red) and predicted values (blue) for hospital and respirator prevalence. Prevalence data is based on NIPaR and may be different to the data from Helsedirektoratet.

2 National 3-week predictions: Hospital beds and Ventilator bed

In this section we show the projected national prevalence of hospitalised patients (hospital beds) and prevalence of patients receiving ventilator treatment (ventilator beds). The projected epidemic and healthcare burden are illustrated in table 2.

Age-specific hospital prevalence predictions are provided in Figures 4 and 5.

Table 2: Estimated national hospital beds and ventilator beds. Median/Mean (CI)

	1 week prediction (Jun 26)	2 week prediction (Jul 03)	3 week prediction (Jul 10)
Hospital beds	101/96 (47-187)	109/102 (47-214)	116/106 (47-245)
Ventilator beds	5/4 (1-10)	5/5 (1-12)	6/5 (1-13)




Figure 3: National 3 week predictions for hospital beds (left) and ventilator beds (right)




Figure 4: Simulated hospital prevalence by age group taking into account the omicron takeover. Real data is shown as black dots.




Figure 5: Simulated respirator prevalence by age group taking into account the omicron takeover. Real data is shown as black dots.

3 Estimated regional reproduction numbers

Calibration of our regional SMC model to hospitalisation incidence data leads to the following estimates for current regional effective reproduction numbers by county (Table 3).

Below we show the estimated daily number of COVID-19 patients admitted to hospital in each county. Model estimates are shown with blue medians and interquantile bands, which are compared to the actual true data, provided in red. The blue bands describe the uncertainty in the calibrated parameters, in addition to the stochastic elements of our model. Last four data points are shown in black as they may be affected by reporting delay.

Table 3: Regional estimates, 01 Jun-04 Jun

County	Median	95CI	Prob>1
Oslo	0.52	0.29-0.88	0.01
Rogaland	0.39	0.16-0.77	0
Møre og Romsdal	0.81	0.48-1.37	0.21
Nordland	1.07	0.59-1.94	0.59
Viken	0.91	0.58-1.52	0.36
Innlandet	0.91	0.49-1.69	0.37
Vestfold og Telemark	1.32	0.75-2.24	0.83
Agder	0.89	0.46-1.65	0.35
Vestland	0.81	0.47-1.51	0.25
Trøndelag	1.13	0.7-1.96	0.69
Troms og Finnmark	1.06	0.58-1.92	0.58










Figure 6: The map shows the direction of the trend in incidence in the counties based on the latest effective reproduction numbers shown in the other chart. The trend is increasing if the probability that the latest reproduction number is above one is above 95%, the trend is likely increasing if this probability is between 80% and 95%, the trend is uncertain if the probability is between 20% and 80%, the trend is likely decreasing if the probability is between 5% and 20% and is decreasing if the probability that the latest R is above one is less than 5%.

Estimated vs observed hospital incidence and 3 weeks forecast by county:

Forecasts are now based on the estimated reproduction numbers obtained by our regional SMC model, for each county. In the forecasted period of three weeks, we use the reproductive numbers showed on Table 3.





4 Regional 3-week predictions: Cumulative (total) incidence and Prevalence

Below is shown the daily incidence (table 4) and prevalence (table 5) for each county.

Table 4: Predicted incidence per day: Median/Mean (CI)

Region	1 week prediction (27 Jun)	2 weeks prediction (04 Jul)	3 weeks prediction (11 Jul)
Agder	65/164 (9-956)	70/227 (8-1516)	76/287 (8-2218)
Innlandet	112/191 (29-861)	126/260 (29-1391)	142/337 (28-2084)
Møre og Romsdal	49/90 (9-419)	57/125 (9-672)	66/168 (9-970)
Nordland	86/227 (11-1414)	104/336 (11-2502)	125/443 (10-3445)
Oslo	244/349 (88-1302)	286/476 (90-2166)	334/592 (89-2609)
Rogaland	13/18 (2-60)	14/20 (2-72)	14/23 (3-91)
Troms og Finnmark	79/186 (11-1034)	98/288 (11-1916)	121/401 (11-3052)
Trøndelag	312/793 (50-5042)	369/1101 (47-8019)	427/1313 (45-8659)
Vestfold og Telemark	490/1178 (62-6844)	656/1661 (61-9475)	837/1924 (63-8339)
Vestland	154/412 (26-2666)	145/506 (24-3694)	144/578 (23-4720)
Viken	543/918 (152-4067)	578/1148 (147-5885)	612/1366 (145-7872)

Table 5: Predicted prevalence. Number of infectious individuals (asymptomatic plus pre-symptomatic plus symptomatic) per day. Median/Mean and 95 perc. CI for three weeks prediction.

Region	27 Jun	04 Jul	11 Jul	low CI, 11 Jul	high CI, 11 Jul
Agder	299/687	316/953	342/1221	57	9126
Innlandet	499/807	557/1092	625/1424	149	8395
Møre og Romsdal	234/397	267/539	308/720	68	4000
Nordland	380/932	454/1385	549/1865	68	14324
Oslo	1454/1853	1636/2389	1834/2915	832	11639
Rogaland	110/130	111/139	113/152	65	440
Troms og Finnmark	356/765	436/1186	532/1675	76	12252
Trøndelag	1369/3247	1602/4597	1863/5629	264	37932
Vestfold og Telemark	2058/4725	2779/6888	3552/8278	327	37457
Vestland	751/1794	719/2200	699/2543	171	19881
Viken	2627/4111	2783/5100	2938/6074	923	32980

5 Regional 3-week predictions: Hospital beds and ventilator beds

Below is shown the estimated short-term forecasting of expected hospital prevalence (table 6) and patients on ventilator treatment for each county (table 7). These estimates do not take into account imported cases from abroad that may be admitted to hospital.

Table 6: Number of hospitalisation beds occupied by Covid-19 patients: Median/Mean (CI)

Region	1 week prediction (27 Jun)	2 weeks prediction (04 Jul)	3 weeks prediction (11 Jul)
Agder	3/7 (0-34)	3/9 (0-56)	4/13 (0-87)
Innlandet	6/9 (1-34)	6/11 (0-55)	7/15 (0-84)
Møre og Romsdal	3/4 (0-19)	3/6 (0-29)	3/8 (0-42)
Nordland	4/9 (0-49)	5/14 (0-94)	6/20 (0-151)
Oslo	13/16 (3-49)	15/22 (4-85)	18/30 (4-135)
Rogaland	1/1 (0-5)	1/1 (0-5)	1/1 (0-6)
Troms og Finnmark	4/7 (0-37)	5/12 (0-69)	6/18 (0-127)
Trøndelag	15/29 (1-154)	17/44 (2-298)	21/60 (2-417)
Vestfold og Telemark	20/41 (2-210)	27/66 (2-391)	36/90 (2-480)
Vestland	9/19 (1-107)	9/24 (1-152)	8/29 (0-211)
Viken	26/39 (6-152)	29/49 (5-225)	31/62 (6-331)

Table 7: Number of ICU beds occupied by Covid-19 patients: Median/Mean (CI)

Region	1 week prediction (27 Jun)	2 weeks prediction (04 Jul)	3 weeks prediction (11 Jul)
Agder	0/0 (0-2)	0/0 (0-3)	0/1 (0-4)
Innlandet	0/0 (0-2)	0/1 (0-3)	0/1 (0-4)
Møre og Romsdal	0/0 (0-1)	0/0 (0-2)	0/0 (0-2)
Nordland	0/0 (0-2)	0/1 (0-4)	0/1 (0-6)
Oslo	0/0 (0-2)	0/1 (0-3)	0/1 (0-4)
Rogaland	0/0 (0-1)	0/0 (0-1)	0/0 (0-1)
Troms og Finnmark	0/0 (0-2)	0/0 (0-3)	0/1 (0-5)
Trøndelag	1/1 (0-6)	1/2 (0-11)	1/3 (0-18)
Vestfold og Telemark	1/1 (0-7)	1/2 (0-15)	2/4 (0-22)
Vestland	0/1 (0-5)	0/1 (0-7)	0/1 (0-9)
Viken	1/2 (0-7)	2/2 (0-9)	2/3 (0-14)

6 14-day trend analysis of confirmed cases and hospitalisations by county



To estimate recent trends in hospitalisation and number of positive tests, we present results in table 8 based on a negative binomial regression where we account for weekend effects. We exclude the last three days to avoid problems of reporting delay and fit the model using data from 17 days to 3 days before the current date. We fit a separate trend model for confirmed cases and for hospital incidence. We only fit a trend model if there has been more than 5 cases or hospitalisations in the 14-day period.

Table 8: Trend analysis for the last 14 days


County	Average daily increase last 14 days		Doubling Time (days)	
	Hospitalisations	Cases	Hospitalisations	Cases
Agder	1.7 (-11.4, 17) %	4.9 (0.1, 10) %	40.8 (-5.7, 4.4)	14.5 (1142.4, 7.3)
Innlandet	-21.6 (-36.5, -7.5) %	-2.9 (-7.5, 1.8) %	-2.8 (-1.5, -8.9)	-23.2 (-8.9, 38.7)
Møre og Romsdal	19.3 (2.6, 42.5) %	5.9 (1.9, 10.1) %	3.9 (26.9, 2)	12.1 (37.8, 7.2)
Nordland	14.3 (3, 28.1) %	8.4 (4.1, 13) %	5.2 (23.2, 2.8)	8.6 (17.2, 5.7)
Norge	4 (0.6, 7.5) %	6.2 (2.4, 10.2) %	17.7 (117.9, 9.5)	11.5 (29.6, 7.1)
Oslo	3.9 (-5, 13.9) %	5.2 (2.1, 8.3) %	18 (-13.6, 5.3)	13.8 (33.2, 8.7)
Rogaland	Not enough data	12.3 (5.2, 20) %	Not enough data	6 (13.8, 3.8)
Troms og Finnmark	1.9 (-13, 19.9) %	4.7 (-2.8, 12.9) %	36.1 (-5, 3.8)	15.2 (-24.3, 5.7)
Trøndelag	8.7 (-0.9, 19.8) %	9.2 (3.2, 15.7) %	8.3 (-78.9, 3.8)	7.9 (22.1, 4.8)
Vestfold og Telemark	5.8 (-2.3, 14.9) %	7.5 (1.6, 13.7) %	12.3 (-29.4, 5)	9.6 (44, 5.4)
Vestland	-7.8 (-15.6, 0.3) %	6.3 (2.2, 10.5) %	-8.5 (-4.1, 228)	11.4 (31.6, 6.9)
Viken	6.5 (0.7, 12.9) %	5.6 (0, 11.5) %	11 (105.9, 5.7)	12.7 (1683.6, 6.4)

7 Regional SMC-model: Estimated daily reproduction numbers


In the figures below we plot the 95% credibility interval and quantiles of the estimated posterior distribution of the regional, daily reproduction numbers. For some counties, uncertainty is large towards the most recent time, because there are very few data and possibly reporting delays which are different in each county.




Vestfold og Telemark






Agder



Rogaland







8 Short-term predictions per health trust ("helseforetak"): number of occupied covid-19 beds and number of patients receiving ventilator treatment

In the table below we distribute our short-term predictions to the Norwegian health trusts ("helseforetak"), on the basis of their areas of responsibility.

Table 9: Number of hospitalisation beds occupied by Covid-19 patients: Median/Mean (CI)


Helseforetak	1 week prediction (27 Jun)	2 weeks prediction (04 Jul)	3 weeks prediction (11 Jul)
Akershus universitetssykehus	13/17 (4-58)	14/22 (5-88)	16/28 (5-128)
Finnmarkssykehuset	1/2 (0-11)	2/4 (0-21)	2/6 (0-40)
Helse Bergen	7/14 (1-77)	6/17 (1-107)	6/20 (0-151)
Helse Fonna	1/2 (0-12)	1/3 (0-18)	1/4 (0-25)
Helse Førde	2/3 (0-18)	2/4 (0-26)	2/5 (0-36)
Helse Møre og Romsdal	3/4 (0-18)	3/6 (0-27)	3/8 (0-42)
Helse Nord Trøndelag	4/9 (1-48)	5/14 (1-95)	7/19 (1-132)
Helse Stavanger	1/1 (0-3)	1/1 (0-3)	1/1 (0-4)
Nordlandssykehuset	2/5 (0-29)	3/8 (0-53)	3/11 (0-88)
Oslo universitetssykehus	11/13 (3-40)	13/19 (3-70)	15/25 (3-111)
St. Olavs hospital	10/20 (1-104)	12/30 (1-202)	14/41 (1-283)
Sykehuset Innlandet	6/8 (1-30)	6/11 (1-48)	7/14 (1-75)
Sykehuset Telemark	8/17 (1-87)	11/27 (1-163)	15/37 (1-198)
Sykehuset i Vestfold	11/24 (1-123)	16/39 (1-231)	21/53 (2-281)
Sykehuset Østfold	7/10 (2-39)	7/13 (2-58)	8/16 (2-84)
Sørlandet sykehus	3/7 (0-33)	4/9 (0-56)	4/13 (0-86)
Universitetssykehuset i Nord-Norge	3/6 (0-28)	4/9 (0-52)	5/14 (0-94)
Vestre Viken	10/15 (3-59)	11/19 (2-89)	12/24 (2-130)

Table 10: Number of respirator beds occupied by Covid-19 patients: Median/Mean (CI)


Helseforetak	1 week prediction (27 Jun)	2 weeks prediction (04 Jul)	3 weeks prediction (11 Jul)
Akershus universitetssykehus	1/1 (0-2)	1/1 (0-3)	1/1 (0-5)
Finnmarkssykehuset	0/0 (0-1)	0/0 (0-1)	0/0 (0-2)
Helgelandssykehuset	0/0 (0-1)	0/0 (0-1)	0/0 (0-2)
Helse Bergen	0/1 (0-4)	0/1 (0-5)	0/1 (0-6)
Helse Fonna	0/0 (0-1)	0/0 (0-1)	0/0 (0-1)
Helse Førde	0/0 (0-1)	0/0 (0-1)	0/0 (0-2)
Helse Møre og Romsdal	0/0 (0-1)	0/0 (0-2)	0/0 (0-2)
Helse Nord Trøndelag	0/0 (0-2)	0/1 (0-4)	0/1 (0-6)
Helse Stavanger	0/0 (0-1)	0/0 (0-1)	0/0 (0-1)
Nordlandssykehuset	0/0 (0-1)	0/0 (0-2)	0/0 (0-3)
Oslo universitetssykehus	0/0 (0-2)	0/1 (0-3)	0/1 (0-3)
St. Olavs hospital	1/1 (0-4)	1/1 (0-7)	1/2 (0-12)
Sykehuset Innlandet	0/0 (0-2)	0/0 (0-3)	0/1 (0-4)
Sykehuset Telemark	0/1 (0-3)	0/1 (0-6)	1/2 (0-9)
Sykehuset i Vestfold	1/1 (0-4)	1/1 (0-9)	1/2 (0-13)
Sykehuset Østfold	0/0 (0-2)	1/1 (0-2)	1/1 (0-4)
Sørlandet sykehus	0/0 (0-2)	0/0 (0-3)	0/1 (0-4)
Universitetssykehuset i Nord-Norge	0/0 (0-1)	0/0 (0-2)	0/1 (0-4)
Vestre Viken	0/1 (0-3)	1/1 (0-4)	1/1 (0-6)

In the figures below, we plot the 95% credibility interval and quantiles of the estimated posterior distribution of the daily number of occupied covid-19 beds and the number of patients receiving ventilator treatment for a few health trusts. The others are available on request. In blue are actual data, in green our median prediction and in gray the 95% credibility interval. For some counties, uncertainty is considerable towards the most recent time because there is little data and possibly reporting delays that are different in each county.


Akershus universitetssykehus




Akershus universitetssykehus




Oslo universitetssykehus




Oslo universitetssykehus



St. Olavs hospital



St. Olavs hospital



9 Mobility

Number of trips out from each municipality during each day is based on Telenor mobility data. The reference level is set to 100 on March 2nd 2020 for all the figures in this section, and we plot the seven-day, moving average of the daily mobility. Figure 13 shows an overview of the mobility since March 2020 for the largest municipalities in each county, and Figure 14 shows the total mobility out from all municipalities in each county, including Oslo. Figure 15 and 16, zooms in on mobility from August 16 2021, for municipalities and counties, respectively.




Figure 13: Mobility for selected municipalities since March 2020: Nationally (Norge), Stavanger (1103), Ålesund (1507), Bodø (1804), Bærum (3024), Ringsaker (3411), Sandefjord (3804), Kristiansand (4204), Bergen (4601), Trondheim (5001), Tromsø (5401).




Figure 14: Mobility for fylker since March 2020: Oslo (03), Rogaland (11), Møre og Romsdal (15), Nordland (18), Viken (30), Innlandet (34), Vestfold og Telemark (38), Agder (42), Vestland (46), Trøndelag (50), Troms og Finmark (54).




Figure 15: Zoom: Mobility from August 16, 2021 and onwards: Nationally (Norge), Stavanger (1103), Ålesund (1507), Bodø (1804), Bærum (3024), Ringsaker (3411), Sandefjord (3804), Kristiansand (4204), Bergen (4601), Trondheim (5001), Tromsø (5401).




Figure 16: Zoom: Mobility from August 16, 2021 and onwards, per fylker: Oslo (03), Rogaland (11), Møre og Romsdal (15), Nordland (18), Viken (30), Innlandet (34), Vestfold og Telemark (38), Agder (42), Vestland (46), Trøndelag (50), Troms og Finnmark (54).

	21	22	23	24	25
Norge	86.9	82.7	84.1	87.9	89.5
Stavanger	82.3	79.4	73.0	78.5	87.5
Ålesund	91.5	88.9	90.2	93.4	100.1
Bodø	94.3	91.4	101.0	100.3	112.9
Bærum	67.4	67.3	58.3	60.8	65.9
Ringsaker	77.8	75.5	75.3	79.4	78.0
Sandefjord	76.9	71.3	73.5	75.4	73.1
Kristiansand	98.7	88.5	98.3	99.4	103.8
Bergen	88.8	86.4	85.4	88.8	93.6
Trondheim	101.2	96.6	97.9	102.5	114.7
Tromsø	108.1	112.2	127.2	125.4	141.6

Table 11: Municipalities

	21	22	23	24	25
Oslo	73.4	68.8	64.8	68.5	76.0
Rogaland	82.9	79.8	75.0	80.3	84.7
Møre og Romsdal	98.5	95.6	96.3	101.5	102.5
Nordland	104.9	106.0	121.7	116.8	124.1
Viken	79.4	73.8	72.2	75.4	77.5
Innlandet	92.6	89.1	93.5	98.7	97.5
Vestfold og Telemark	86.5	81.1	86.5	90.2	86.4
Agder	98.4	90.6	101.6	104.0	100.9
Vestlandet	93.2	90.5	94.4	99.7	96.4
Trøndelag	102.8	99.7	102.1	108.5	112.6
Troms og Finnmark	103.4	102.4	116.6	115.6	122.7

Table 12: Counties

Weekly mobility for Norway and selected municipalities is displayed in Table 11 and mobility for counties is displayed in Table 12. The percentages in the tables are to be interpreted towards the reference level of 100 for week 10 in March 2020. The color-coding encodes the following: 'Green' monotonic decrease in mobility, 'Yellow' almost monotonic decrease or flat mobility trend, 'Red' increasing mobility.

9.1 Foreign roamers on Telenor's network in Norway

9.1 Foreign roamers on Telenor's network in Norway

An analysis of foreign roamers in Norway from January 2020 has been carried out, to better understand the potential virus importation. In Figure 17 the total number of roamers per day per county are displayed.

Figure 18 shows the levels of roamers from the following countries: Poland, Lithuania, Sweden, Netherlands, Denmark, Latvia, Germany, Spain, Finland and the rest of the world. These levels represent the total number of foreign, visiting roamers from each of the countries per day in Norway, since July 5 2021.




Figure 17: The total number of foreign roamers in Norway broken down on different fylker: Oslo (3), Rogaland (11), Møre og Romsdal (15), Nordland (18), Viken (30), Innlandet (34), Vestfold og Telemark (38), Agder (42), Vestland (46), Trøndelag (50), Troms og Finnmark (54).

9.1 Foreign roamers on Telenor's network in Norway











Figure 18: National overview of total number of foreign, visiting roamers from Poland, Lithuania, Sweden, Netherlands, Denmark, Latvia, Germany, Spain, Finland and the rest of the world.

9.2 Foreign roamers per county (fylke) in Norway

9.2 Foreign roamers per county (fylke) in Norway



9.2 Foreign roamers per county (fylke) in Norway



10 Estimated parameters

Table 13: Estimated parameters

	Min.	1st Qu.	Median	Mean	3rd Qu.	Max.	Period
R0s	1.945	2.512	2.694	2.701	2.875	3.626	Until March 14
R1s	0.471	0.637	0.685	0.683	0.727	0.883	From 15 March to 19 April
R2s	0	0.313	0.49	0.487	0.641	1.297	From 20 April to 10 May
R3s	0.005	0.272	0.525	0.545	0.757	1.631	From 11 May to 30 June
R4s	0.004	0.321	0.563	0.581	0.778	1.696	From 01 July to 31 July
R5s	0.201	1.065	1.212	1.2	1.351	1.726	From 01 August to 31 August
R6s	0.719	0.943	1.007	1.003	1.066	1.354	From 01 September to 25 October
R7s	0.574	0.924	1.107	1.13	1.31	2.075	From 26 October to 04 November
R8s	0.741	0.931	0.984	0.98	1.034	1.18	From 05 November to 30 November
R9s	0.674	0.812	0.859	0.855	0.901	1.051	From 01 December to 03 January
R10s	0.387	0.66	0.783	0.779	0.899	1.259	From 04 January to 21 January
R11s	0.397	0.75	0.859	0.878	1.01	1.541	From 22 January to 07 February
R12s	0.968	1.211	1.291	1.295	1.387	1.697	From 08 February to 01 March
R13s	0.808	0.959	1.006	1.007	1.053	1.215	From 02 March to 24 March
R14s	0.698	0.811	0.837	0.838	0.866	0.944	From 25 March to 05 May
R15s	0.563	0.803	0.886	0.884	0.965	1.33	From 06 May to 26 May
R16s	0.222	0.616	0.73	0.729	0.835	1.281	From 27 May to 20 June
R17s	0.14	0.668	0.8	0.781	0.913	1.245	From 21 June to 04 August
R18s	0.791	1.03	1.092	1.092	1.157	1.354	From 05 August to 31 August
R19s	0.507	0.698	0.753	0.754	0.81	1.026	From 01 September to 24 September
R20s	0.927	0.998	1.021	1.02	1.042	1.122	From 25 September to 14 December
R21s	0.915	1.032	1.066	1.068	1.106	1.233	From 15 December to 13 January
R22s	1.255	1.313	1.336	1.337	1.36	1.426	From 14 January to 25 February
R23s	0.853	0.909	0.928	0.928	0.947	1.006	From 26 February to 15 May
R24s	0.804	1.136	1.228	1.219	1.309	1.599	From 16 May

Table 14: Assumptions

Assumptions	Mean	Distribution	Reference
Mobile Mobility Data			
Telenor coverage	48%		https://ekomstatistikken.nkom.no/
Data updated	Jun 5		
Data used in the predictions	Jun 3 rd	Fixed	Corrected to preserve population
Model parameters			
Exposed period ($1/\lambda_1$)	2 days	Exponential	changed from Feretti et al 2020
Pre-symptomatic period ($1/\lambda_2$)	2 days	Exponential	Feretti et al 2020
Symptomatic infectious period ($1/\gamma$)	3 days	Exponential	changed from Feretti et al 2020
Asymptomatic, infectious period ($1/\gamma$)	3 days	Exponential	changed from Feretti et al 2020
Infectiousness asympt. (r_{I_a})	0.1	Fixed	Feretti et al 2020
Infectiousness presymp (r_{E_2})	1.3	Fixed	guided by Feretti et al 2020
Prob. asymptomatic infection (p_a)	0.4		Feretti et al 2020
Healthcare			
Fraction asymptomatic infections	40%	Fixed	Mizumoto et al 2020 20% for the old population, Diamond Princess
% symptomatic and asymptomatic infections requiring hospitalization:			Salje et al 2020 corrected for: % of elderly living in elderly homes in Norway (last two age groups) and corrected for presence among positive tested since May 1.
0-9 years	0.1%	Fixed	
10 - 19 years	0.1%		
20 - 29 years	0.5%		
30 - 39 years	1.1%		
40 - 49 years	1.4%		
50 - 59 years	2.9%		
60 - 69 years	5.8%		
70 - 79 years	9.3%		
80+ years	22.3%		
Probability that an admission has been reported on Monday			
From Sunday	32%	Fixed	Estimated from "Beredskapsregistret BeredtC19"
From Saturday	49%		
From Friday	68%		
From Thursday	86%		
Probability that an admission has been reported			
From one day before	53%	Fixed	Estimated from "Beredskapsregistret BeredtC19"
From two days before	77%		
From three days before	82%		
From four days before	91%		
Probability that a positive laboratory test has been reported			
From one day before	6.7%	Fixed	Estimated from MSIS
From two days before	59%		
From three days before	90%		
From four days before	97%		
Probability that a negative laboratory test has been reported			
From one day before	16%	Fixed	Estimated from MSIS
From two days before	74%		
From three days before	92%		
From four days before	98%		

Supplementary analysis: Trend analysis using various data sources

Due to significant changes in test recommendations, we present trend calculations from several monitoring data sources. Seen together, they can give insights on the trend of new infections. The data sources include:

- MSIS - A trend calculated from the number of confirmed positive cases. Changes in test criteria in recent weeks will likely lead to underestimating the trend of new infections.
- NOPaR: Admissions with covid-19 as the main reason, the trend is calculated from daily numbers admissions with covid-19 as the main cause. It usually takes longer from infection to hospitalization than for testing; therefore, this indicator lags behind the trend of transmission at around 1-2 weeks. The major difference in severity between Delta and Omicron also makes this indicator more challenging to interpret in the transition phase between the two variants.
- Symptometer - From the Symptometer survey, we calculate the proportion of respondents who self-report symptoms and a positive test for covid-19. This indicator provides a measure of the prevalence of infection in society, but after conversion, it can also provide an estimate of a trend for new infections.
- sKUHR - We use the development of the number of consultations with confirmed covid-19 (R992) at GP offices and emergency rooms, adjusting for delayed reporting the last 14 days. After correction for the delay between onset and seeking a doctor, this indicator measures the incidence of infection.
- MoBa - The participants in MoBa are sent a mobile questionnaire form every other week, including questions about who has been "sick with respiratory symptoms/fever the last 14 days" and how many days ago the symptoms started. We calculate an approximate 14-day incidence of symptom onset from this indicator, which is used to estimate a trend in infection. The number of respondents is about 60-75,000 in each round.
- NPR and NoPaR: Proportion of all acute admissions with covid-19, including entries with other main causes than covid-19. This indicator measures the prevalence of infection in society and is then converted into incidence of infection.




Figure 21: Trend analysis using various data sources




Figure 22: Regional hospitalisation probabilities per infection. The estimates are based on Salje et al., and regional data on the age distribution in the test data and the empirical case-hospitalisation rates.

Models and materials:

This report presents results based on a mathematical infectious disease model describing the geographical spread of COVID-19 in Norway. We use a metapopulation model for situational awareness and short-term forecasting and an individual-based model for long-term predictions. This report does not contain the long-term prediction results. Reproduction numbers of the metapopulation model are estimated in two ways: SMC-ABC is used to estimate a step-function in the transmissibility through prespecified changepoints, and SMC is used to estimate a daily varying reproduction number. We also provide estimates based on EpiEstim and a simple trend analysis. The models are described in previous reports and will not be explained here.

The metapopulation model takes daily varying Telenor mobility data as input. We also provide plots of the recent mobility for situational awareness.

How you should interpret the results: 3-week-ahead predictions and long-term scenarios

We provide both 3-week-ahead predictions and long-term scenarios. These are simulations of the disease spread into the future, under specific assumptions.

In the 3-week-ahead predictions, we assume that all parameters are as today, and simulate disease spread 3-weeks-ahead in time. Hence, these predictions are conditional on the current situation, and specifically on the most recently estimated reproduction number. The 3-week-ahead predictions thus do not take into account changes in transmissibility that are not yet captured by the available data, for example due to the delay between transmission and hospitalisation. Hence one of the conditions for the predictions to be valid is that the intervention policies do not change significantly in the next weeks. Hence, it does not make sense to evaluate or use the predictions if there are big changes in factors like

- new interventions
- relaxation of interventions
- a combination of new interventions and relaxations
- a significant change in vaccination coverage
- new variants with new properties
- a significant change in the contact behaviour of individuals.

As these factors are not likely to stay constant in the long-term future, we do not produce predictions for longer than three weeks ahead in time. Hence, our 3-week-ahead predictions are predictions of what may happen in the future, if there were no significant changes in the assumptions.

In this report, the term patient in ventilator treatment includes only those patients that require either invasive mechanical ventilation or ECMO (Extracorporeal membrane oxygenation).

FHI COVID-19 modelling team:

- **Birgitte Freiesleben de Blasio** - Department of Method Development and Analytics. Norwegian Institute of Public Health and Oslo Centre for Biostatistics and Epidemiology, University of Oslo.
- **Francesco Di Ruscio** - Department of Method Development and Analytics. Norwegian Institute of Public Health.
- **Gunnar Øyvind Isaksson Rø** - Department of Method Development and Analytics. Norwegian Institute of Public Health.
- **Solveig Engebretsen** - Norsk Regnesentral.
- **Arnoldo Frigessi** - Oslo Centre for Biostatistics and Epidemiology, University of Oslo and Oslo University Hospital.
- **Alfonso Diz-Lois Palomares** - Department of Method Development and Analytics. Norwegian Institute of Public Health.
- **Magnus Nygård Osnes** - Department of Method Development and Analytics. Norwegian Institute of Public Health.
- **Anja Bråthen Kristoffersen** - Department of Method Development and Analytics. Norwegian Institute of Public Health.
- **Kenth Engø-Monsen** - Telenor Research.
- **Louis Yat Hin Chan** - Department of Method Development and Analytics. Norwegian Institute of Public Health.
- **Jonas Christoffer Lindstrøm** - Department of Method Development and Analytics. Norwegian Institute of Public Health.
- **Richard White** - Department of Method Development and Analytics. Norwegian Institute of Public Health.
- **Gry Marysol Grøneng** - Department of Method Development and Analytics. Norwegian Institute of Public Health.
- **Chi Zhang** - Department of Method Development and Analytics. Norwegian Institute of Public Health.
- **Jørgen Eriksson Midtbø** - Department of Method Development and Analytics. Norwegian Institute of Public Health.
- **Geir Storvik** - Department of Mathematics. University of Oslo.